

Azimuthally-sensitive interferometry and the source lifetime at RHIC

Michael A. Lisa

for the STAR Collaboration

Abstract Pion interferometry (HBT) measurements relative to the reaction plane provide an estimate of the transverse source anisotropy at freeze-out, which probes the system dynamics and evolution duration. Measurements by the STAR Collaboration indicate that the source is extended increasingly out-of-plane with increasing impact parameter, suggesting a short evolution duration roughly consistent with estimates based on azimuthally-integrated HBT measurements.

Key words interferometry • HBT • heavy ion collisions • freeze-out

Introduction

Two-particle intensity interferometry (Hanbury-Brown-Twiss, or HBT) analyses of relativistic heavy ion collisions represent the most direct experimental probes of the space-time structure of the source generated in such collisions [15]. At the Relativistic Heavy Ion Collider (RHIC), identical-pion HBT studies in Au+Au collisions [5, 7, 21] yielded an apparent source size quantitatively consistent with measurements at lower energies, in contrast to predictions of larger sources based on QGP formation [26, 27]. In addition, hydrodynamical models, successful at RHIC in describing momentum-space quantities such as transverse momentum spectra and elliptic flow [6], have failed to reproduce the small HBT radii [16]. This HBT puzzle [11, 12] may arise because the system's lifetime is shorter than predicted by models.

In non-central collisions, the initial anisotropic collision geometry generates greater transverse pressure gradients in the reaction plane than perpendicular to it. This leads to preferential in-plane expansion (elliptic flow) [6, 19, 22, 31] which diminishes the initial spatial anisotropy. The final (freeze-out) source shape should be sensitive to the evolution of the pressure gradients and the system lifetime; a long-lived source would be less out-of-plane extended and perhaps in-plane extended. Hydrodynamic calculations [18] predict a strong sensitivity of the HBT parameters to the early source conditions and show that, while the source may still be out-of-plane extended after hydrodynamic evolution, a subsequent rescattering phase [30] tends to make the source in-plane. Thus, the freeze-out source shape might discriminate between scenarios of the system's evolution.

Here, we present results of a measurement of azimuthally-sensitive HBT in Au+Au collisions at $\sqrt{s_{NN}} = 200$ GeV. We discuss how these results may be used to

M. A. Lisa
Physics Department,
The Ohio State University,
Columbus, OH 43210, USA,
Tel.: +614 292 8524, Fax: +614 292 7557,
e-mail: lisa@mps.ohio-state.edu

Received: 28 January 2004

extract, in a model-dependent way, the geometric anisotropy of the source at freeze-out. Finally, we use this extracted geometry to estimate the evolution timescale of the system.

Measurement

The measurements were made using the STAR detector [2] at RHIC. Particle trajectories and momenta were reconstructed using a Time Projection Chamber (TPC) with full azimuthal coverage, located inside a 0.5 Tesla solenoidal magnet. Au+Au event samples of 0–5%, 5–10%, 10–20%, 20–30%, and 30–80% of the total cross hadronic section (based on charged particle multiplicity) are presented here. The second-order event plane angle Ψ_2 [23] for each event was determined from the weighted sum of primary charged-particle transverse momenta (\bar{p}_T) [1]. Within the resolution which we determine from the random subevent method [23], $\Psi_2 \approx \Psi_{rp}$ (true reaction plane angle) or $\Psi_2 \approx \Psi_{rp} + \pi$; i.e. the direction of the impact parameter vector is determined up to a sign [23], and the measured event plane is roughly coplanar with the true reaction plane [4, 24].

Pions were selected according to their specific energy loss (dE/dx) in the TPC in the rapidity range $|y| < 0.5$. Pair-wise cuts to remove two-track merging and single-track splitting effects were applied to signal and background distributions [7].

Pairs of like-sign pions were placed into bins of $\Phi = \phi_{pair} - \Psi_2$, where ϕ_{pair} is the azimuthal angle of the total pair momentum ($\mathbf{k} = \mathbf{p}_1 + \mathbf{p}_2$). Because we use the 2nd-order

event plane, Φ is only defined in the range $(0, \pi)$. For each bin, a three-dimensional correlation function is constructed in the “out-side-long” decomposition [9] of the relative momentum \mathbf{q} .

Finite reaction plane resolution and finite width of the Φ bins has the effect of reducing the measured oscillation amplitudes. A model-independent correction procedure [14] is applied to each \mathbf{q} -bin in the numerator and denominator of each correlation function, the overall effect of which is to increase the amplitude of the oscillations of the HBT radii vs. Φ (~ 10 –30%).

To account for the effect of final-state Coulomb interactions, we adopt the approach first proposed by Bowler [10] and Sinyukov [29] and recently advocated by the CERES collaboration [3]. We fit each experimental correlation function to the form:

$$(1) \quad C(q, \Phi) = N \cdot [(1 - \lambda) \cdot 1 + \lambda \cdot K(q)(1 + G(q, \Phi))]$$

where the $(1 - \lambda)$ and λ terms account for the non-participating and participating fractions of pairs, respectively, $K(Q_{inv})$ is the square of the Coulomb wave-function, N is the normalization, and $G(q, \Phi)$ is the Gaussian form [9]:

$$(2) \quad G(q, \Phi) = e^{-q_0^2 R_0^2(\Phi) - q_s^2 R_s^2(\Phi) - q_l^2 R_l^2(\Phi) - q_o q_s R_{os}^2(\Phi)}$$

R_i^2 the squared HBT radii, where the l, s, o subscripts indicate the long (parallel to beam), side (perpendicular to beam and total pair momentum) and out (perpendicular to q_l and q_s) decomposition of q .




Fig. 1. Squared HBT radii measured relative to the reaction plane, for three selections of event centrality.




Fig. 2. Blast-wave [25] calculations of Fourier coefficients for the oscillations of squared HBT radii, for three sources of varying transverse eccentricity. Arrows indicate approximation of Eq. (4).

Results, Fourier coefficients and estimating eccentricity

Figure 1 shows the squared HBT radii as a function of Φ , for three centrality classes and $0.15 \leq k_T \leq 0.6$ GeV/c. Curves correspond to a Fourier decomposition of the squared-radii oscillations according to [13, 14]

$$(3) \quad R_{\mu,n}^2(k_T) = \begin{cases} \langle R_{\mu}^2(k_T, \Phi) \cos(n\Phi) \rangle (\mu = o, s, l) \\ \langle R_{\mu}^2(k_T, \Phi) \sin(n\Phi) \rangle (\mu = o, s) \end{cases}$$

As expected [7], the 0th-order Fourier coefficients (FCs), i.e. the mean value of the radii, indicate larger source sizes for more central collisions. We verified that the 0th-order FC for, R_o^2 , R_s^2 and R_l^2 correspond to HBT radii obtained in an azimuthally-integrated analysis. Higher-order ($n > 2$) FCs were found to be negligible.

HBT radii directly measure homogeneity regions [15, 28] which constitute only part of the whole source; hence,

connections to the overall source geometry inevitably rely on a model. The so-called blast-wave model [25] has been relatively successful in describing soft physics observables (spectra, elliptic flow, azimuthally-integrated HBT) at RHIC. In the context of a generalized blast-wave, with spatial and flow anisotropies, a recent study [25] finds that the transverse source shape is related to $R_s^2(\Phi)$ as

$$(4) \quad \varepsilon \equiv \frac{(R_y^2 - R_x^2)}{(R_y^2 + R_x^2)} \approx \frac{1}{2} \cdot \frac{R_{s,2}^2}{R_{s,0}^2} \approx \frac{1}{2} \cdot \frac{R_{o,2}^2}{R_{s,0}^2} \approx \frac{1}{2} \cdot \frac{R_{os,2}^2}{R_{s,0}^2}$$

where R_x and R_y are the source length scales in and out of the reaction plane, respectively. The approximation becomes exact in the case of vanishing transverse flow [20, 32]. In Fig. 2 are the 0th and 2nd-order FCs, as defined in Eq. (3) for three sources as calculated in the blast-wave. The sources have the same temperature, radial flow, timescales, and average size (all adjusted roughly to the values which




Fig. 3. Fourier coefficients for the oscillations of squared HBT radii, for 3 k_T bins, vs. centrality (number of participating nucleons) measured by STAR. Left panels: means (0th-order FC) of oscillations; right panels: relative 2nd-order oscillation amplitudes. See text for details.

best describe RHIC data), but differing ϵ . Changing from a round ($\epsilon = 0$) to a deformed source increases the oscillations of the transverse radii (and, very slightly – note the scale) the longitudinal radius. Eq. (4) arises from the observation that 2ϵ , indicated by small arrows on the right panels, approximately reproduces the relative oscillations in the transverse radii. Therefore, relying on the blast-wave model for context, we may estimate the freeze-out source shape.

Figure 3 shows the measured FCs, as a function of centrality, for three bins in k_T , quantifying the increasing size (left panels) and decreasing anisotropy (right panels) as the collision becomes more central. In Fig. 4, we compare the initial anisotropy, $\epsilon_{initial}$, calculated with a Glauber model outlined in Ref. [8] with ϵ_{final} , estimated by applying Eq. (4) to the data of Fig. 3. We note that ϵ_{final} has a meaningful sign, and its positive values indicate that the timescales and/or flow velocities of these collisions are not sufficiently large to fully quench the initial out-of-plane geometric anisotropy of the system.

A connection to timescales

The data in Fig. 3 should provide useful constraints on timescales and evolution in transport models of heavy ion collisions. As a crude example, here we take the model-

dependent data in Fig. 4 one step further. Motivated by hydrodynamical calculations [17], we assume that the transverse flow strength grows roughly linearly with time, and anisotropies in the flow field set in very early. Since there is no collective flow initially, the flow velocity and source extensions in (x) and out of (y) the reaction plane evolve with time as




Fig. 4. Final (from HBT) vs. initial (from Glauber) transverse source eccentricity, estimated from the data in Fig. 3 and Eq. (4). The line represents $\epsilon_{initial} = \epsilon_{final}$.




Fig. 5. Toy-model calculations for the evolution of the transverse eccentricity as a function of time. See text for details.

$$(5) \quad \beta_{x,y}(t) = \frac{t}{\tau_0} \cdot \beta_{x,y}^{FO} \rightarrow R_{x,y}(t) = R_{x,y}^{\text{Glauber}} + \frac{1}{2} \left(\frac{\beta_{x,y}^{FO}}{\tau_0} \right) t^2.$$

Here, β^{FO} is the flow velocity (at the edge of the source) at freeze-out, and τ_0 is the evolution time of the system – from initial overlap until freeze-out. From Eq. (5), we may calculate $\epsilon(t)$ in this toy model.

For collisions between 10–20% of the cross-section (3rd centrality bin in Figs. 3 and 4), the estimated initial and final anisotropies are indicated in Fig. 5. Estimates for were taken from blast-wave fits [25] to mid-central collisions at $\sqrt{s_{NN}} = 130$ GeV. In order to evolve ϵ , we must assume a value for τ_0 . Blast-wave fits [25] to azimuthally-integrated longitudinal HBT radii suggest a rather short evolution time of ~ 9 fm/c. Assuming this value, we arrive at a consistent picture: as shown by the solid line in Fig. 5, at $t = \tau_0 = 9$ fm/c, [epsilon] has reached its measured value. On the other hand, using a significantly different value for τ_0 yields inconsistent results, as shown by the dashed and dotted lines. Thus, our measurements, in the context of this admittedly crude model, seem to provide corroborating evidence in the direction of evolution timescales on the order of 9 fm/c.

Summary

We have presented an analysis of azimuthally-sensitive pion interferometry for Au+Au collisions at RHIC. We observe clear 2nd-order oscillations of the transverse HBT radii, and extract Fourier coefficients. Geometrically, as the impact parameter decreases, the 0th-order FCs increase, indicating larger sources, and the relative oscillations decrease, suggesting rounder ones. More quantitative statements rely on model interpretations.

In the blast-wave model, the relative oscillations provide a good estimate of the source ellipticity at freeze-out. In

this context, we find that the source at all centralities is extended out of the reaction plane (though not as much as the initial source), suggesting rather short evolution durations.

This data should provide strong constraints on dynamic models of heavy ion collisions, particularly as regards timescales and pressure anisotropies. To illustrate the point, we performed a toy model calculation which turned out to favor the rather short evolution timescales extracted via other means previously.

References

1. Ackermann K, Adams N, Adler C *et al.* (2001) Elliptic flow in Au+Au collisions at $s(\text{NN})^{1/2} = 130$ GeV. *Phys Rev Lett* 86:402–407
2. Ackermann KH, Adams N, Adler C (for the STAR Collaboration) (2003) STAR detector overview. *Nucl Instrum Meth Phys Res A* 499:624–632
3. Adamova D, Agakichiev G, Appelshauer H (for the CERES Collaboration) (2003) Beam energy and centrality dependence of two-pion Bose-Einstein correlations at SPS energies. *Nucl Phys A* 714:124–144
4. Adams J, Adler C, Aggarwal MM (for the STAR Collaboration) (2004) Azimuthal anisotropy at RHIC: the first and fourth harmonics. (nucl-ex/0310029)
5. Adcox K, Adler SS, Ajitanand NN (for the PHENIX Collaboration) (2002) Centrality dependence of $+/-, K+/-, p$, and production from $s(\text{NN})^{1/2} = 130$ GeV Au+Au collisions at RHIC. *Phys Rev Lett* 88:242301
6. Adler C, Ahammed Z, Allgower C (for the STAR Collaboration) (2001) Identified particle elliptic flow in Au+Au collisions at $s(\text{NN})^{1/2} = 130$ GeV. *Phys Rev Lett* 87:182301
7. Adler C, Ahammed Z, Allgower C *et al.* (for the STAR Collaboration) (2001) Pion interferometry of $s(\text{NN})^{1/2} = 130$ GeV Au+Au collisions at RHIC. *Phys Rev Lett* 87:082301
8. Adler C, Ahammed Z, Allgower C (2002) Centrality dependence of high hadron suppression in Au+Au collisions at $s(\text{NN})^{1/2} = 130$ GeV. *Phys Rev Lett* 89:202301
9. Bertsch G, Gong M, Tohyama M (1988) Pion interferometry in ultrarelativistic heavy-ion collisions. *Phys Rev C* 37:1896–1900
10. Bowler MG (1991) Coulomb corrections to Bose-Einstein correlations have been greatly exaggerated. *Phys Lett B* 270:69–74
11. Dumitru A (2002) How the HBT-puzzle at RHIC might dissipate. (nucl-th/0206011)
12. Heinz U (2003) The quark-gluon plasma at RHIC. *Nucl Phys A* 721:c30–c39
13. Heinz U (2004) Emission angle dependence of HBT radii: Theoretical background and interpretation. (nucl-th/0311048)
14. Heinz U, Hummel A, Lisa MA, Wiedemann UA (2002) Symmetry constraints for the emission angle dependence of Hanbury-Brown-Twiss radii. *Phys Rev C* 66:044903
15. Heinz U, Jacak BV (1999) Two-particle correlations in relativistic heavy-ion collisions. *Annu Rev Nucl Part Sci* 49:529–579
16. Heinz U, Kolb PF (2002) Early thermalization at RHIC. *Nucl Phys A* 702:269–280
17. Kolb PF (2003) A small, but sensitive observable for heavy ion collisions. *Phys Rev C* 68:031902
18. Kolb PF, Heinz U (2003) Emission angle dependent HBT at RHIC and beyond. *Nucl Phys A* 715:653–656
19. Kolb PF, Sollfrank J, Heinz U (2000) Anisotropic transverse flow and the quark-hadron phase transition. *Phys Rev C* 62:054909

20. Lisa MA, Heinz U, Wiedemann UA (2000) Tilted pion sources from azimuthally sensitive HBT interferometry. *Phys Lett B* 489:287–292
21. López Noriega M (for the STAR Collaboration) (2003) Pion interferometry in Au Au collisions at $s(\text{NN})^{1/2} = 200$ GeV. *Nucl Phys A* 715:623–626
22. Ollitrault JY (1997) Reconstructing azimuthal distributions in nucleus-nucleus collisions. (nucl-ex/9711003)
23. Poskanzer AM, Voloshin SA (1998) Methods for analyzing anisotropic flow in relativistic nuclear collisions. *Phys Rev C* 58:1671–1678
24. Pratt S, Csörgő T, Zimanyi J (1990) Detailed predictions for two-pion correlations in ultrarelativistic heavy-ion collisions. *Phys Rev C* 42:2646–2652
25. Retière F, Lisa MA (2003) Observable implications of geometrical and dynamical aspects of freeze-out in heavy ion collisions. (nucl-th/0312024)
26. Rischke DH (1996) Hydrodynamics and collective behaviour in relativistic nuclear collisions. *Nucl Phys A* 610:88–101
27. Rischke DH, Gyulassy M (1996) The time-delay signature of quark-gluon plasma formation in relativistic nuclear collisions. *Nucl Phys A* 608:479–512
28. Sinyukov YuM (1994) Spectra and correlations in locally equilibrium hadron and quark-gluon systems. *Nucl Phys A* 566:589–592
29. Sinyukov YuM, Lednický R, Akkelin SV, Pluta J, Erazmus B (1998) Coulomb corrections for interferometry analysis of expanding hadron systems. *Phys Lett B* 432:248–257
30. Teaney D, Lauret J, Shuryak E (2001) A hydrodynamic description of heavy ion collisions at the SPS and RHIC. (nucl-th/0110037)
31. Voloshin SA (2003) Anisotropic flow. *Nucl Phys A* 715:379–388
32. Wiedemann UA (1998) Two-particle interferometry for noncentral heavy-ion collisions. *Phys Rev C* 57:266–279